


# What can COVID-19 shutdowns teach us about reducing air pollution?

The impact of COVID-19 on manufacturing, travel and the associated pollutant production has proved an unexpected opportunity for a global experiment for atmospheric scientists. Drs Jenny Stavrakou and Maite Bauwens at the Royal Belgian Institute for Space Aeronomy are experts in atmospheric modelling and interpretation of satellite measurements of air pollution. They have been exploring how COVID-related changes have affected our environment and how we might use this to improve air quality in the future.



The atmosphere is a complicated place. Thousands of chemical reactions take place unceasingly, with a constant flux of new species coming in and out. Some of those reactions are triggered by different chemicals colliding with each other. Others happen when molecules absorb light from the sun.

The sheer number of different reactions and possible reaction routes makes modelling this chemical soup very challenging. Different layers of the atmosphere are home to different types of chemistry as not all wavelengths of light can pass through all the layers. Different regions tend to vary in their chemical composition as well.


Despite its complexity, scientists internationally are working very hard to understand how our atmosphere is formed, as its rich chemistry has a direct impact on the quality of the air we breathe. Air pollution, harmful gases and particles in the air are serious global problems and are thought to be responsible for around 7 million deaths worldwide each year.

Of these air pollutants, nitrogen dioxide (NO<sub>2</sub>) is particularly

detrimental in terms of impact on human health and the environment. High concentrations of NO<sub>2</sub> itself can irritate the lungs and aggravate other respiratory diseases, like asthma. NO<sub>2</sub> can also go on to react and form particulate matter and ozone – both of which have similar, problematic effects. In the environment, NO<sub>2</sub> contributes to acid rain, damaging forests, coasts and even causing buildings in our cities to crumble.

Dr Jenny Stavrakou and Dr Maite Bauwens at the Royal Belgian Institute for Space Aeronomy have been tracking NO<sub>2</sub> levels using a special

Antuna/Shutterstock.com



tool, the Tropospheric Monitoring Instrument (TROPOMI), aboard the Sentinel-5 Precursor satellite mission, dedicated to the daily monitoring of air pollution worldwide. They have been watching with great interest how NO<sub>2</sub> emissions have been strongly reduced in response to the COVID-related restrictions of human activity, e.g. stay-at-home orders, travel bans and a fall in manufacturing. From the data they are recording, the team is working out exactly what the changing patterns of NO<sub>2</sub> emissions mean for our environment and atmosphere, and what we can learn to help reduce air pollution in the future.

**NO<sub>2</sub> SOURCES**  
Generally, NO<sub>2</sub> and related



Sentinel-5 Precursor.

molecules (known as NO<sub>x</sub>) are formed predominantly from the combustion of fossil fuels. Motor vehicles are a common source and even oil or gas burning at home, such as gas cookers, produce NO<sub>2</sub>. While

**NO<sub>2</sub> itself can irritate the lungs and aggravate other respiratory diseases.**

very low concentrations are harmless, it can start to have negative effects when levels exceed 0.1 ppm.

The constant circulation and mixing of the atmosphere mean that even if the pollution is created in a localised area, it will eventually circulate globally. With TROPOMI, Drs Stavrakou and Bauwens have been focusing on the day-to-day change in NO<sub>2</sub> levels in cities. While they have found some reduction in NO<sub>2</sub> levels from new environmental legislation aimed at minimising air pollution, the most significant change has arisen from the COVID-19 pandemic.

What the researchers learned was that, in many of the epicentres of the pandemic, the reduction in human activity caused a sharp and dramatic reduction in NO<sub>2</sub> levels.

Drs Stavrakou and Bauwens could therefore take a temporary glimpse of a world without nearly as much NO<sub>2</sub>.

**WINDOW OF OPPORTUNITY**  
Unfortunately for the environment,

NO<sub>2</sub> levels largely rebounded as economic activities have restarted, particularly in China and India. While global economic activities are still somewhat reduced post-pandemic, this has given Drs Stavrakou and Bauwens unique data to use to try and understand exactly what and how different factors contributed to the observed reduction in NO<sub>2</sub> levels. While reduced economic activity played a role, meteorological factors also influence NO<sub>2</sub> levels.


By having such extensive daily measurements on NO<sub>2</sub> levels, the data from TROPOMI can be used to refine atmospheric models and gives us a chance to see whether such models are still robust and reliable with a sudden reduction in the number of NO<sub>2</sub> sources globally.

TROPOMI, which was launched on the Sentinel-5 satellite back in 2017, is not the only instrument to measure our atmosphere. Drs Stavrakou and Bauwens have also been comparing this with data from the Ozone Monitoring Instrument (OMI), another satellite launched in 2004, and found an excellent consistency between the two satellite datasets.



Lemberg Vector studio/Shutterstock.com





Average TROPOMI NO<sub>2</sub> column in 2019 (left) and 2020 (right) for the first 2 months after the lockdown measures. The last column shows the absolute column reduction in 10<sup>15</sup> molec. cm<sup>-2</sup>.

#### PROBLEM GASES

One of the reasons NO<sub>2</sub> and the whole family of nitrogen oxides are so problematic for health and the environment are their effects on ozone. There is some concern that NO<sub>2</sub> and ozone can work together to enhance their negative effects on the respiratory system. In the stratosphere, nitrogen oxides can destroy ozone, leading to a thinning of the global ozone layer which protects us from dangerous solar ultraviolet light.

an unprecedented drop in emissions in many regions around the world and cast light on the response of the atmospheric system to the emission disruption caused by the pandemic. Generally, regions with reduced NO<sub>x</sub> production also showed decreased ozone production, but this was not the case everywhere. Surprisingly, in heavily polluted regions during winter, surface ozone has showed increases in response to the drop of NO<sub>2</sub> pollution. This is because the

wide and large the knock-on effects from NO<sub>x</sub> could be.

#### AIR QUALITY CONTROL

As lockdowns occurred in the world during different time periods with different degrees of severity (which could be mapped in how much the NO<sub>2</sub> emissions were reduced) a wealth of data was generated showing how smaller or bigger NO<sub>2</sub> concentration reductions would impact the overall atmospheric chemical composition and

how all of this could also be influenced by the weather conditions.

As TROPOMI keeps on measuring NO<sub>x</sub> emission levels, COVID-19


lockdowns have provided a unique chance to examine how human activity has influenced the composition of our atmosphere, with many cities enjoying periods of significantly reduced pollution hazes and less respiratory irritation. Now, the question remains how to use this new knowledge to inform public policy going forward to address another public health crisis – air pollution.

**The reduction of NO<sub>x</sub> concentrations during the pandemic also led to a cascading change in the concentrations of many other chemical species.**

Drs Stavrakou and Bauwens's measurements make full use of the high-resolution capabilities of TROPOMI which allow them to identify exactly which regions and areas the NO<sub>2</sub> emission originates from. Their measurements showed a reduction of up to 50% in NO<sub>x</sub> emission in some of China's cities while COVID-19 lockdowns were in place. They have also been able to compare their satellite observations to local surface measurements. Models were able to simulate for the first time

relationship between NO<sub>2</sub> and ozone concentrations and overall air quality is highly nonlinear. This shows that a reduction of NO<sub>2</sub> emissions is often not enough to solve the pollution problem as a whole. The reduction of NO<sub>x</sub> concentrations during the pandemic also led to a cascading change in the concentrations of many other chemical species, showing how

# Behind the Research



**Dr Jenny Stavrakou**

E: [jenny@aeronomie.be](mailto:jenny@aeronomie.be)  
T: +32 (0)2 3736766



**Dr Maite Bauwens**

E: [maite.bauwens@aeronomie.be](mailto:maite.bauwens@aeronomie.be)  
T: +32 (0)2 89 09 877

W: <http://tropo.aeronomie.be>

## Research Objectives

Drs Stavrakou and Bauwens observed the change in NO<sub>2</sub> levels in major cities during COVID-19 shutdowns and analysed the evolution of other major pollutants. Their objective is to interpret these observations using atmospheric models and to infer emissions estimates from space using sophisticated inversion techniques.

## Detail

Royal Belgian Institute for Space Aeronomy (BIRA-IASB)  
Avenue Circulaire 3, 1180 Brussels, Belgium

#### Bio

**Dr Jenny Stavrakou** graduated with a PhD in Physics at the University of Aix-Marseille (France) and joined the research group of Tropospheric Chemistry Modelling at BIRA-IASB in 2001, where she now occupies a Senior Scientist position.

**Dr Maite Bauwens** is postdoctoral researcher at the group of Tropospheric Chemistry Modelling of BIRA-IASB. In 2010 she received her PhD in Sciences from the University of Brussels (VUB), Belgium.

#### Funding

- Belgian Science Policy Office (BELSPO) through the PRODEX TROVA project (2016-present)
- European Space Agency (ESA) through the ICOVAC project

#### Collaborators

- Jean-François Müller, Head of the Tropospheric Chemistry Department, for his valuable advice
- Karolien Lefever, Head of the Communication Department, for her help, discussions and support throughout this study
- All staff of the communication cell who have contributed and are still contributing to this project
- ESA Sentinel-5P Mission team for making TROPOMI satellite data publicly available
- NO<sub>2</sub> satellite product development team from the KNMI institute in the Netherlands and the German Aerospace Center (DLR)

## References

Bauwens, M., Compennolle, S., Stavrakou, T., Müller, J.-F. et al. (2020). Impact of Coronavirus Outbreak on NO<sub>2</sub> Pollution Assessed Using TROPOMI and OMI Observations. *Geophysical Research Letters*, 47(11), e2020GL087978. Available at: <https://doi.org/10.1029/2020GL087978>

Shi, X. and Brasseur, G.P. (2020). The response in air quality to the reduction of Chinese Economic Activities during the COVID-19 Outbreak. *Geophysical Research Letters*, 47(11), e2020GL088070. Available at: <https://doi.org/10.1029/2020GL088070>

Gaubert, B., Bouarar, I., Doumbia, T., Liu, Y. et al. (2020). Global Changes in Secondary Atmospheric Pollutants during the 2020 COVID-19 Pandemic. *Journal of Geophysical Research: Atmosphere*. Available at: <https://doi.org/10.1002/essoar.10504703.1> [Preprint]

## Personal Response

**Do you think the COVID-19 pandemic has raised awareness of air pollution and the importance of minimising NO<sub>2</sub> emissions?**

/// The air pollution changes due to the lockdowns attracted a great deal of media attention and mobilised in an unprecedented way the community of atmospheric scientists to work together and analyse the observed air pollution patterns. Besides its tragic dimension, the pandemic has shown that the air pollution problem is crucial and must be urgently tackled. As clean air swiftly returned in many polluted cities, from New York to Delhi and Beijing, the pandemic offered a glimpse of what can be achieved by speeding up the implementation of sustainable policies to make our cities healthier. ///

